

American Journal of Physiology: Renal, Fluid and Electrolyte Physiology

No. 1. JANUARY 1994

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Structural determinants of glomerular hydraulic permeability <i>M. C. Drumond and W. M. Deen</i> | F1 |
| Calcium and free radicals in hypoxia/reoxygenation injury of renal epithelial cells <i>E. L. Greene and M. S. Paller</i> | F13 |
| Functional and cytoskeletal changes induced by sublethal injury in proximal tubular epithelial cells <i>V. M. Kroshian, A. M. Sheridan, and W. Lieberthal</i> | F21 |
| Effect of dietary sodium chloride on insulin receptor number and mRNA levels in rat kidney <i>L. A. Sechi, C. A. Griffin, and M. Schambelan</i> | F31 |
| Regulation of prostaglandin endoperoxide synthase gene expression in rat mesangial cells by interleukin-1 β <i>D. Rzymkiewicz, K. Leingang, N. Baird, and A. R. Morrison</i> | F39 |
| Mechanism of endothelin activation of phospholipase A ₂ in rat renal medullary interstitial cells <i>R. L. Barnett, L. Ruffini, D. Hart, P. Mancuso, and E. P. Nord</i> | F46 |
| Postnatal maturation of potassium transport in rabbit cortical collecting duct <i>L. M. Satlin</i> | F57 |
| Interaction of rat kidney P-glycoprotein with a urinary component and various drugs including cyclosporin A <i>J. H. M. Charuk, T. W. Loo, D. M. Clarke, and R. A. F. Reithmeier</i> | F66 |
| Mineralocorticoid receptors and 11 β -steroid dehydrogenase activity in renal principal and intercalated cells <i>A. N  ray-Fejes-T  th, E. Rusvai, and G. Fejes-T  th</i> | F76 |
| Glomerular endothelial cells in culture express and secrete vascular endothelial growth factor <i>K. Uchida, S. Uchida, K. Nitta, W. Yumura, F. Marumo, and H. Nihei</i> | F81 |
| Angiotensin II suppression is a major factor permitting excretion of an acute sodium load in humans <i>D. R. J. Singer, N. D. Markandu, J. J. Morton, M. A. Miller, G. A. Sagnella, and G. A. MacGregor</i> | F89 |
| Intracellular modulation of acid secretion in rat inner medullary collecting duct cells <i>J. H. Schwartz, S. A. Masino, R. D. Nichols, and E. A. Alexander</i> | F94 |
| Insulin-like growth factor I produces renal hyperfiltration by a kinin-mediated mechanism <i>A. A. Jaffa, D. LeRoith, C. T. Roberts, Jr., P. F. Rust, and R. K. Mayfield</i> | F102 |
| Effects of corticoid agonists and antagonists on apical Na ⁺ permeability of toad urinary bladder <i>H. Garty, K. Peterson-Yantorno, C. Asher, and M. M. Civan</i> | F108 |
| Natriuretic response to renal interstitial hydrostatic pressure during angiotensin II blockade <i>J. A. Haas, J. C. Lockhart, T. S. Larson, T. Henrikson, and F. G. Knox</i> | F117 |
| Augmentation of intrarenal angiotensin II levels by chronic angiotensin II infusion <i>A. M. Von Thun, R. C. Vari, S. S. El-Dahr, and L. G. Navar</i> | F120 |
| Hepatocyte growth factor accelerates recovery from acute ischemic renal injury in rats <i>S. B. Miller, D. R. Martin, J. Kissane, and M. R. Hammerman</i> | F129 |
| Effect of an endothelin-receptor antagonist on ischemic acute renal failure <i>L. Chan, A. Chittinandana, J. I. Shapiro, P. F. Shanley, and R. W. Schrier</i> | F135 |
| Vasopressin-independent regulation of collecting duct water permeability <i>J. S. Han, Y. Maeda, C. Ecelbarger, and M. A. Knepper</i> | F139 |

- IGF binding protein 5 and IGF-I receptor regulation in hypophysectomized rat kidneys
*M. K. Hise, N. M. Mantzouris, J. S. Lahn, M. S. Sheikh, Z.-M. Shao,
 and J. A. Fontana* F147

RAPID COMMUNICATION

- Spatially restricted expression of *set* mRNA in developing rat kidney
E.-G. Kim, M. E. Choi, and B. J. Ballermann F155

No. 2. FEBRUARY 1994

EDITORIAL REVIEW

- Hereditary and acquired defects in signaling through the hormone-receptor-G
 protein complex
J. R. Raymond F163

BRIEF REVIEW

- Renal cortical interstitium and fluid absorption by peritubular capillaries
K. Aukland, R. T. Bogusky, and E. M. Renkin F175

- Metabolic acidosis stimulates carbonic anhydrase activity in rabbit proximal tubule and
 medullary collecting duct
L. P. Brion, B. J. Zavilowitz, C. Suarez, and G. J. Schwartz F185

- Ca²⁺ uptake, fatty acid, and LDH release during proximal tubule hypoxia: effects of
 mepacrine and dibucaine
*D. Bunnachak, A. R. P. Almeida, J. F. M. Wetzels, P. Gengaro, R. A. Nemenoff,
 T. J. Burke, and R. W. Schrier* F196

- Effect of luminal angiotensin II on proximal tubule fluid transport: role of apical
 phospholipase A₂
L. Li, Y.-P. Wang, A. W. Capparelli, O. D. Jo, and N. Yanagawa F202

- ATP stimulates Ca²⁺ mobilization by a nucleotide receptor in glomerular endothelial cells
V. A. Briner and F. Kern F210

- Mechanism of acidification along cortical distal tubule of the rat
*R. Fernandez, M. J. Lopes, R. F. de Lira, W. F. G. Dantas, E. J. Cragoe, Jr.,
 and G. Malnic* F218

- Synergistic effects of angiotensin and adenosine in the renal microvasculature
H. Weihprecht, J. N. Lorenz, J. P. Briggs, and J. Schnermann F227

- Expression of Na-K-ATPase α - and β -subunit mRNA and protein isoforms in the
 rat nephron
J. A. Tumlin, C. A. Hoban, R. M. Medford, and J. M. Sands F240

- Intracellular processing of immune complexes formed on the surface of glomerular
 epithelial cells
A. K. Singh and M. A. Rahman F246

- Effect of U-73,122, an inhibitor of phospholipase C, on actions of parathyroid hormone in
 opossum kidney cells
*K. J. Martin, C. L. McConkey, A. K. Jacob, E. A. Gonzalez, M. Khan,
 and J. J. Baldassare* F254

- Conductive properties of papillary surface epithelium
W. B. Reeves F259

- Polarized distribution of oxalate transport systems in LLC-PK₁ cells, a line of renal
 epithelial cells
H. Koul, S. Ebisuno, L. Renzulli, M. Yanagawa, M. Menon, and C. Scheid F266

- Inhibition of renal vascular 20-HETE production impairs autoregulation of renal
 blood flow
*A.-P. Zou, J. D. Imig, M. Kaldunski, P. R. Ortiz de Montellano, Z. Sui,
 and R. J. Roman* F275

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Molecular adaptations of GLUT1 and GLUT2 in renal proximal tubules of diabetic rats <i>J. H. Dominguez, K. Camp, L. Maianu, H. Feister, and W. T. Garvey</i> | F283 |
| Endothelin-1 production by rat inner medullary collecting duct: effect of nitric oxide, cGMP, and immune cytokines <i>D. E. Kohan and E. Padilla</i> | F291 |
| Growth and characterization of porcine urinary bladder epithelial cells in vitro <i>C. Guhe and W. Föllmann</i> | F298 |
| X-linked hypophosphatemic Gy mice: renal tubular maximum for phosphate vs. brush-border transport after low-P diet <i>S. W. Thornton, H. S. Tenenhouse, J. Martel, R. W. Bockian, M. H. Meyer, and R. A. Meyer, Jr.</i> | F309 |
| Regulation of rat renal α_2B -adrenergic receptors by potassium depletion <i>J. C. Muir, L. Huang, J. K. Harrison, D. L. Rosin, and M. D. Okusa</i> | F316 |
| Ionic currents and endothelin signaling in smooth muscle cells from rat renal resistance arteries <i>D. V. Gordienko, C. Clausen, and M. S. Goligorsky</i> | F325 |

RAPID COMMUNICATION

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Characterization of human uroguanylin: a member of the guanylin peptide family <i>T. Kita, C. E. Smith, K. F. Fok, K. L. Duffin, W. M. Moore, P. J. Karabatsos, J. F. Kachur, F. K. Hamra, N. V. Pidhorodeckyj, L. R. Forte, and M. G. Currie</i> | F342 |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|

LETTERS TO THE EDITOR

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Effect of angiotensin II on renal arteriolar resistances estimated by morphometric analysis <i>K. Endlich and M. Steinhausen; W. P. Anderson, K. M. Denton, and D. Alcorn</i> | F349 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|

ANNOUNCEMENTS

F352

No. 3. MARCH 1994

EDITORIAL REVIEW

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Why renal cysts grow <i>K. D. Gardner, Jr., R. H. Glew, A. P. Evan, J. A. McAteer, and J. Bernstein</i> | F353 |
| Erythropoietin enhances recovery from cisplatin-induced acute renal failure <i>N. D. Vaziri, X. J. Zhou, and S. Y. Liao</i> | F360 |
| pH _i determines rate of sodium transport in frog skin: results of a new method to determine pH _i <i>R. Rick</i> | F367 |
| Glycogen metabolism in dog inner medullary collecting ducts <i>L. Meury, J. Sénécal, J. Noël, and P. Vinay</i> | F375 |
| Differential regulation of angiotensin II and losartan binding sites in glomeruli and mesangial cells <i>D. Chansel, T. Bizet, S. Vandermeersch, P. Pham, B. Levy, and R. Ardaillou</i> | F384 |
| Effect of insulin on water and urea transport in the inner medullary collecting duct <i>A. J. Magaldi, K. R. César, and Y. Yano</i> | F394 |
| cAMP stimulates proximal convoluted tubule Na ⁺ -K ⁺ -ATPase activity <i>S. Breton, J. S. Beck, and R. Laprade</i> | F400 |
| Systemic and renal effect of intravenous infusion of endothelin-1 in healthy human volunteers <i>S. S. Sørensen, J. K. Madsen, and E. B. Pedersen</i> | F411 |

| | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Collecting duct changes in potassium depletion: effects of ACE inhibition <i>S. K. Mujais</i> | F419 |
| Mechanisms for protective effects of free radical scavengers on gentamicin-mediated nephropathy in rats <i>T. Nakajima, A. Hishida, and A. Kato</i> | F425 |
| Metabolic responses to nephrosis: effect of a low-protein diet <i>E.-J. Choi, J. Bailey, R. C. May, T. Masud, and B. J. Maroni</i> | F432 |
| Basolateral taurine transport system in reptilian renal cells <i>S. Benyajati and S. M. Bay</i> | F439 |
| Peritubular organic cation transport in isolated rabbit proximal tubules <i>C. E. Groves, K. K. Evans, W. H. Dantzler, and S. H. Wright</i> | F450 |
| Effects of epidermal growth factor and transforming growth factor- α on rabbit proximal tubule solute transport <i>R. Quigley and M. Baum</i> | F459 |
| Mechanisms of bicarbonate transport by cultured rabbit inner medullary collecting duct cells <i>A. E. Weill, C. C. Tisher, M. F. Conde, and I. D. Weiner</i> | F466 |
| Immunolocalization of calcitriol receptor, 24-hydroxylase cytochrome P-450, and calbindin D _{28k} in human kidney <i>R. Kumar, J. Schaefer, J. P. Grande, and P. C. Roche</i> | F477 |
| Increased cytosolic Ca ²⁺ inhibits AVP-stimulated adenylyl cyclase activity in rat IMCT cells by activation of PKC <i>I. Teitelbaum and T. Berl</i> | F486 |
| Synthesis and localization of C-type natriuretic peptide in mammalian kidney <i>A. D. Dean, V. M. Vehaskari, and J. E. Greenwald</i> | F491 |
| Parathyroid hormone inhibits Na ⁺ -K ⁺ -ATPase through a cytochrome P-450 pathway <i>C. M. P. Ribeiro, G. R. Dubay, J. R. Falck, and L. J. Mandel</i> | F497 |
| Arginine vasopressin and forskolin regulate apical cell surface expression of epithelial Na ⁺ channels in A6 cells <i>T. R. Kleyman, S. A. Ernst, and B. Coupaye-Gerard</i> | F506 |

RAPID COMMUNICATION

| | |
|-----------------------------------------------------------------------------------------------------------------------------|------|
| Bicarbonate dependency of betaine synthesis in cultured LLC-PK ₁ cells <i>G. W. Moeckel and Y.-H. H. Lien</i> | F512 |
|-----------------------------------------------------------------------------------------------------------------------------|------|

ANNOUNCEMENTS

F516

No. 4. APRIL 1994

EDITORIAL REVIEW

| | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Coupling between transepithelial Na transport and basolateral K conductance in renal proximal tubule <i>J. S. Beck, R. Laprade, and J.-Y. Lapointe</i> | F517 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|------|

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Cellular actions of cAMP on HCO ₃ ⁻ -secreting cells of rabbit CCD: dependence on in vivo acid-base status <i>C. Emmons and J. B. Stokes</i> | F528 |
| Na pump defects in chronic uremia cannot be attributed to changes in Na-K-ATPase mRNA or protein <i>S. Greiber, B. K. England, S. R. Price, R. M. Medford, R. G. Ebb, and W. E. Mitch</i> | F536 |

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Chloride channels in apical membrane of primary cultures of rabbit distal bright convoluted tubule <i>V. Poncet, M. Tauc, M. Bidet, and P. Poujeol</i> | F543 |
| Trafficking of apical proteins into clathrin-coated vesicles isolated from rat renal cortex <i>T. G. Hammond and P. J. Verroust</i> | F554 |
| Feeding-induced increase in urinary dopamine excretion is independent of renal innervation and sodium intake <i>B. Mühlbauer and H. Osswald</i> | F563 |
| Macrophage Fc receptor activity modulates mesangial cell proliferation and matrix synthesis <i>J. Mattana and P. C. Singhal</i> | F568 |
| Long-term effects of growth factors on pH and acid-base transport in rat glomerular mesangial cells <i>M. B. Ganz and W. F. Boron</i> | F576 |
| Modification of mesangial cell function by chloride is attenuated in spontaneously hypertensive rats <i>T. Okuda, Y. Inishi, T. Arakawa, M. Ohara, and K. Kurokawa</i> | F586 |
| Combined K ⁺ and Cl ⁻ repletion corrects augmented H ⁺ secretion by distal tubules in chronic alkalosis <i>D. E. Wesson</i> | F592 |
| Renin release from permeabilized juxtaglomerular cells is stimulated by chloride but not by low calcium <i>B. L. Jensen and O. Skøtt</i> | F604 |
| Glyceraldehyde-3-phosphate dehydrogenase modifier protein is associated with microtubules in kidney epithelial cells <i>N. H. Aithal, M. M. Walsh-Reitz, S. Kartha, M. P. Janulis, T. E. Martin, and F. G. Toback</i> | F612 |
| Effects of pH on apical calcium entry and active calcium transport in rabbit cortical collecting system <i>R. J. M. Bindels, A. Hartog, S. L. Abrahamse, and C. H. van Os</i> | F620 |
| Early increase in pulsatile growth hormone release after unilateral nephrectomy in adult rats <i>A. Haramati, M. D. Lumpkin, and S. E. Mulroney</i> | F628 |
| Activation of acid-secreting intercalated cells in rabbit collecting duct with ammonium chloride loading <i>J. W. Verlander, K. M. Madsen, J. K. Cannon, and C. C. Tisher</i> | F633 |
| Expression of the D ₂ subfamily of dopamine receptor genes in kidney <i>D.-Q. Gao, L. M. Canessa, M. M. Mouradian, and P. A. Jose</i> | F646 |
| An Arg-Gly-Asp peptide stimulates Ca ²⁺ efflux from osteoclast precursors through a novel mechanism <i>K. Yamakawa, R. Duncan, and K. A. Hruska</i> | F651 |
| Effects of endothelin on rat renal proximal tubule Na ⁺ -P _i cotransport and Na ⁺ /H ⁺ exchange <i>J. Guntupalli and T. D. DuBose, Jr.</i> | F658 |
| Second messengers involved in the control of renin secretion in cultured human nephroblastoma cells <i>A. M. Devlin and B. J. Leckie</i> | F667 |
| Interleukin-1 inhibits sodium and water transport in rabbit cortical collecting duct <i>Y. Sakairi, Y. Ando, K. Tabei, E. Kusano, and Y. Asano</i> | F674 |

SPECIAL COMMUNICATION

| | |
|-------------------------------------------------------------------------------------|------|
| A simplified device for injection of paraffin wax blockades <i>T. L. Pallone</i> | F681 |
|-------------------------------------------------------------------------------------|------|

EDITORIAL REVIEW

- Potential-induced changes in intracellular pH
V. Lyall and T. U. L. Biber F685
- Sodium excess aggravates hypertension and renal parenchymal injury in rats with chronic NO inhibition
C. K. Fujihara, S. M. Michellazzo, G. de Nucci, and R. Zatz F697
- Impaired homologous upregulation of vitamin D receptor in rats with chronic renal failure
H. Koyama, Y. Nishizawa, M. Inaba, M. Hino, J. M. Prahl, H. F. DeLuca, and H. Morii F706
- Interleukin-1 α stimulates KC synthesis in rat mesangial cells: glucocorticoids inhibit KC induction by IL-1
L. Feng, Y. Xia, J. I. Kreisberg, and C. B. Wilson F713
- Long-term regulation of collecting duct water permeability: freeze-fracture analysis of isolated perfused tubules
J. B. Wade, S. Nielsen, R. A. Coleman, and M. A. Knepper F723
- Renal function in hypertensive rats transgenic for mouse renin gene
J. E. Springate, L. G. Feld, and D. Ganten F731
- Role of renal nerves for the expression of renin in adult rat kidney
S. Holmer, B. Rinne, K.-U. Eckardt, M. Le Hir, K. Schrickler, B. Kaissling, G. Riegger, and A. Kurtz F738
- Glomerular response to acute protein load is not blunted by high-protein diet or nephron reduction
M. Burtin, D. Laouari, C. Kindermans, and C. Kleinknecht F746
- Protein restriction sequentially induces new urea transport processes in rat initial IMCD
T. Iozaki, A. G. Gillin, C. E. Swanson, and J. M. Sands F756
- Inhibition of nitric oxide causes exaggerated natriuresis in spontaneously hypertensive rats
A. A. Khraibi F762
- Expression of Na-P₁ cotransport in rat kidney: localization by RT-PCR and immunohistochemistry
M. Custer, M. Löttscher, J. Biber, H. Murer, and B. Kaissling F767
- Role of intracellular Ca²⁺ in modulation of tight junction resistance in A6 cells
B. Jovov, S. A. Lewis, W. E. Crowe, J. R. Berg, and N. K. Wills F775
- Inhibition of Na⁺/Ca²⁺ exchange in renal BLMV by IP₃ depends on site of action and direction of Ca²⁺ flux
C. L. Fraser, C. Cummings, and G. Cassafer F785
- Adenosine A₁ receptor-mediated inhibition of vasopressin action in inner medullary collecting duct
R. M. Edwards and W. S. Spielman F791
- Urate transport in brush-border membrane of human kidney
F. Roch-Ramel, D. Werner, and B. Guisan F797
- Lithium reabsorption in perfused loops of Henle: effects of perfusion rate and bumetanide
R. J. Unwin, S. J. Walter, and D. G. Shirley F806

SPECIAL COMMUNICATION

- A novel approach allows identification of K channels in the lateral membrane of rat CCD
W.-H. Wang, C. M. McNicholas, A. S. Segal, and G. Giebisch F813

RAPID COMMUNICATION

- Distinct cellular localization of mRNAs for three subtypes of prostaglandin E receptor in kidney
Y. Sugimoto, T. Namba, R. Shigemoto, M. Negishi, A. Ichikawa, and S. Narumiya F823

EDITORIAL REVIEW

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| The emerging role of transforming growth factor- β in kidney diseases <i>K. Sharma and F. N. Ziyadeh</i> | F829 |
| <hr/> | |
| Exogenous aldosterone antagonizes distal tubular effects of calcium entry blocker felodipine <i>H. W. van Hamersvelt, J. F. M. Wetzels, H. J. Kloke, R. A. P. Koene, and F. T. M. Huysmans</i> | F843 |
| Vasoconstriction of outer medullary vasa recta by angiotensin II is modulated by prostaglandin E ₂ <i>T. L. Pallone</i> | F850 |
| Endogenous renal dopamine production regulates phosphate excretion <i>A. Debska-Slizien, P. Ho, R. Drangova, and A. D. Baines</i> | F858 |
| H ⁺ -ATPases of renal cortical and medullary endosomes are differentially sensitive to Sch-28080 and omeprazole <i>I. Sabolić, D. Brown, J.-M. Verbavatz, and J. Kleinman</i> | F868 |
| In vivo modulation of rat distal tubule net HCO ₃ flux by VIP, isoproterenol, angiotensin II, and ADH <i>D. Z. Levine, M. Iacovitti, S. Buckman, and V. Harrison</i> | F878 |
| Protein restriction reduces transforming growth factor- β and interstitial fibrosis in nephrotic syndrome <i>A. A. Eddy</i> | F884 |
| Endothelin modulates the mitogenic effect of PDGF on glomerular mesangial cells <i>M. Kohno, T. Horio, K. Yokokawa, K. Yasunari, N. Kurihara, and T. Takeda</i> | F894 |
| Inhibition of bicarbonate transport in peanut lectin-positive intercalated cells by a monoclonal antibody <i>G. Fejes-Tóth, A. Náray-Fejes-Tóth, L. M. Satlin, F. M. Mehrgut, and G. J. Schwartz</i> | F901 |
| Angiotensin II type 1 receptor: role in renal growth and gene expression during normal development <i>A. Tufro-McReddie, D. W. Johns, K. M. Geary, H. Dagli, A. D. Everett, R. L. Chevalier, R. M. Carey, and R. A. Gomez</i> | F911 |
| Distal tubule unidirectional HCO ₃ reabsorption in vivo during acute and chronic metabolic alkalosis in the rat <i>D. Z. Levine, M. Iacovitti, S. Buckman, D. Vondorpe, V. Harrison, and S. P. Nadler</i> | F919 |
| Macrophages, monocyte chemoattractant peptide-1, and TGF- β 1 in experimental hydronephrosis <i>J. R. Diamond, D. Kees-Folts, G. Ding, J. E. Frye, and N. C. Restrepo</i> | F926 |
| Effect of P-450 ω -hydroxylase metabolites of arachidonic acid on tubuloglomerular feedback <i>A.-P. Zou, J. D. Imig, P. R. Ortiz de Montellano, Z. Sui, J. R. Falck, and R. J. Roman</i> | F934 |
| Altered renal vascular responses in the aging rat kidney <i>J. E. Tank, J. P. Vora, D. C. Houghton, and S. Anderson</i> | F942 |
| Rat models for clinical use of insulin-like growth factor I in acute renal failure <i>S. B. Miller, D. R. Martin, J. Kissane, and M. R. Hammerman</i> | F949 |
| Endothelin stimulates phosphatidylcholine hydrolysis through both PLC and PLD pathways in mesangial cells <i>E. Baldi, A. Musial, and M. Kester</i> | F957 |
| ATP-dependent transport of tetraethylammonium by endosomes isolated from rat renal cortex <i>J. B. Pritchard, D. B. Sykes, R. Walden, and D. S. Miller</i> | F966 |
| Differentiation of intercalated cells in developing rat kidney: an immunohistochemical study <i>J. Kim, C. C. Tisher, and K. M. Madsen</i> | F977 |

CORRIGENDA

Volume 265, November 1993

Volume 34, November 1993

Pages F660–F669: J. E. Benabe, S. Wang, J. N. Wilcox, and M. Martinez-Maldonado. "Modulation of ANG II receptor and its mRNA in normal rat by low-protein feeding." Page F662, Table 1: for the third group (NP), the value for renal vascular resistance is incorrect and should be $9 \pm 1 \text{ dyn} \cdot \text{s} \cdot \text{cm}^{-5}$; also for the NP group, the plasma renin activity is incorrectly expressed as a negative value and should be $10.1 \pm 1.3 \text{ ng} \cdot \text{ml}^{-1} \cdot \text{h}^{-1}$.

